

International Journal of Modern Physics A
© World Scientific Publishing Company

The Short Range Mechanism of N-N interaction in the Extended Chiral $SU(3)$ Quark Model *

L. R. Dai[†]

Department of Physics, Liaoning Normal University, 116029, Dalian, P. R. China

Z. Y. Zhang, Y. W. Yu

Institute of High Energy Physics, 100039, Beijing, P. R. China

We give the comparisons between the chiral $SU(3)$ quark model and the extended chiral $SU(3)$ quark model. The results show that the phase shifts of NN scattering are very similar. However, the short range mechanisms of nucleon-nucleon interaction are totally different. In the chiral $SU(3)$ quark model, the short range interaction is dominantly from OGE, and in the extended chiral $SU(3)$ quark model, it is dominantly from vector meson exchanges.

Keywords: NN interaction ; Quark Model; Chiral Symmetry.

1. Introduction

The chiral $SU(3)$ quark model ¹ was proposed by generalizing the idea of the $SU(2)$ σ model to the flavor $SU(3)$ case. In this model, the nonet pseudo-scalar meson exchanges and the nonet scalar meson exchanges are considered in describing the medium and long range parts of the interactions, and the one gluon exchange (OGE) potential is still retained to contribute the short range repulsion. By using this model, the energies of the baryon ground states, the N-N scattering phase shifts and the hyperon-nucleon (Y-N) cross sections can be reproduced reasonably. It seems that the repulsive core of the N-N interaction can be explained by the OGE and the quark exchange effect.

Since last few years, Shen et al ², Riska and Glozman ³ applied the quark-chiral field coupling model to study the baryon structure. They found that the chiral field coupling is also important in explaining the structure of baryons. In the work of Riska et al ³, the vector meson coupling was also included to replace OGE. They pointed out the spin-flavor interaction is important in explaining the energy of the Roper resonance and got a comparatively good fit to the baryon spectra. The quark-chiral field coupling model is a big challenge to the Isgur's model ⁴, in which the

*The project is supported by Scientific Research Foundation of Liaoning Education Department (No. 202122028)

[†]dailr@lnnu.edu.cn

2 *L. R. Dai, Z. Y. Zhang, Y. W. Yu*

OGE governs the baryon structure.

On the other hand, in the traditional one boson exchange (OBE) model on baryon level, the N-N short range repulsion comes from vector meson (ρ, ω, K^* and ϕ) exchanges. Some authors^{5,6} also studied short-range NN repulsion as stemming from the Goldstone boson (and ρ -like) exchanges on the quark level. It has been shown that these interactions can substitute traditional OGE mechanism. Naturally, we would like to ask which is the right mechanism for describing the short range interactions? Is it from OGE or from vector meson exchange or both important? This is interesting and challenging. To study this problem, we propose the extended chiral $SU(3)$ quark model⁷ in which the vector meson exchanges are involved.

2. The Extended Chiral $SU(3)$ Quark Model

2.1. The Model

By generalizing the interaction Lagrangian, on the basis of the chiral $SU(3)$ quark model¹, we further added the term of coupling between quark and vector meson field,

$$\mathcal{L}_I^V = -ig_{chv}\bar{\psi}\gamma_\mu\vec{\varphi}_\mu\cdot\vec{\tau}\psi - i\frac{f_{chv}}{2M_p}\bar{\psi}\sigma_{\mu\nu}\partial_\nu\vec{\varphi}_\mu\cdot\vec{\tau}\psi. \quad (1)$$

Therefore, we got the extended chiral $SU(3)$ quark model⁷, the Hamiltonian of the system in this extended model can be written as

$$H = \sum_i T_i - T_G + \sum_{i<j} V_{ij}, \quad (2)$$

$$V_{ij} = V_{ij}^{Conf} + V_{ij}^{OGE} + V_{ij}^{Ch}, \quad (3)$$

$$V_{ij}^{Ch} = \sum_{a=0}^8 V_{ij}^{s_a} + \sum_{a=0}^8 V_{ij}^{ps_a} + \sum_{a=0}^8 V_{ij}^{v_a}. \quad (4)$$

Compared with the chiral $SU(3)$ quark model¹, we have additional term of quark vector field coupling potential $V_{ij}^{v_a}$ which is derived from Lagrangian in the Eq. (1).

2.2. Determine the Parameters

We have two initial input parameters: the harmonic-oscillator width parameter b_u , and the up (down) quark mass. The coupling constant g_{ch} for scalar and pseudo-scalar chiral field coupling is fixed by experimental value. For vector chiral field coupling, the g_{chv} and f_{chv} are the coupling constants for vector coupling and tensor coupling respectively. In the study of nucleon resonance transition coupling to vector meson, Riska et al took the value of $g_{chv} = 3.0$ and neglected the tensor coupling part⁸. From one boson exchange theory on baryon level, we can obtain the two values between quark and baryon levels. All meson masses, m_{ps} , m_s , and

m_v , are taken to be the experimental values, except for σ meson, its mass is treated as an adjustable parameter. The cut-off mass Λ , indicating the chiral symmetry breaking scale is taken to be $\Lambda^{-1} = 0.18 fm$. In the calculation, η and η' mesons are mixed by η_1 and η_8 with the mixing angle $\theta_\eta = -23^\circ$. ω and ϕ are mixed by ω_1 and ω_8 with the mixing angle $\theta_\omega = -54.7^\circ$. After the parameters of chiral fields are fixed, the one gluon exchange coupling constants g_u can be determined by the mass splits between N and Δ . The confinement strength is fixed by the stability conditions of N. The resultant model parameters are tabulated in Table 1.

Table 1. Model parameters and the binding energies of deuteron.

	chiral $SU(3)$ quark model	Extended chiral $SU(3)$ quark model	
		set I	set II
$b_u(fm)$	0.5	0.45	0.45
$g_{nn\pi}$	13.67	13.67	13.67
g_{ch}	2.621	2.621	2.621
g_{chv}	0	2.351	1.972
f_{chv}/g_{chv}	0	0	2/3
$m_\sigma(MeV)$	595	535	547
g_u	0.886	0.293	0.399
$\alpha_s(g_u^2)$	0.785	0.086	0.159
$a_{uu}(MeV/fm^2)$	48.1	48.0	42.9
$B_{deu}(MeV)$	2.13	2.19	2.14

3. Results and discussions

The N-N phase shifts are calculated by solving a Resonating Group Method (RGM) equation in the extended chiral $SU(3)$ quark model. For comparison with results from different models, the results of chiral $SU(3)$ quark model without vector meson exchanges¹ are drawn with short-dashed curves, while long dashed and solid curves are those obtained from the extended chiral $SU(3)$ quark model⁷ with two different sets of parameters (set I and set II). All S,P,D and F partial waves calculated are very similar in both models, and consistent with the experimental results⁷. Here we just discuss two main results:

(i) The phase shifts of S-wave

The results (in Fig.1) show that the 1S_0 phase shifts are obviously improved in the extended chiral $SU(3)$ quark model. The 3S_1 phase shifts of different models are almost the same. To get the right trend of the phase shift versus scattering energy, the size parameter b_u is taken with different values for these two models. $b_u = 0.50 fm$ in the chiral $SU(3)$ quark model, and $b_u = 0.45 fm$ in the extended chiral $SU(3)$ quark model. This means that the bare radius of baryon becomes smaller when more meson clouds are included in the model. This physical picture looks reasonable.


4 *L. R. Dai, Z. Y. Zhang, Y. W. Yu*

Fig. 1. The NN scattering S-Wave phase shifts

(ii) **The coupling constant of OGE**

When the vector meson field coupling is considered, the coupling constant of OGE is greatly reduced by fitting the mass difference between Δ and N . For both set I and II, $\alpha_s < 0.2$, which is much smaller than the value (0.785) of chiral $SU(3)$ quark model (see Table 1). It means that the OGE interaction is rather weak in the extended chiral $SU(3)$ quark model. Instead, the vector meson exchanges play an important role for the short range interaction between two quarks. Hence, mechanisms of the quark-quark short range interactions of these two models are totally different. In the chiral $SU(3)$ quark model, the short range interaction is dominantly from OGE, and in the extended chiral $SU(3)$ quark model, it is dominantly from vector meson exchanges.

These features are reasonable and helpful in better understanding the short range mechanism of the quark-quark interactions.

References

1. Z. Y. Zhang, Y. W. Yu, and L. R. Dai, *HEP & NP* **20**, 393(1996);
Z. Y. Zhang, Y. W. Yu, P. N. Shen, L. R. Dai, A. Faessler and U. Straub, *Nucl. Phys.* **A625**, 59(1997).
2. P.N. Shen, Y. B. Dong, Y. W. Yu, Z. Y. Zhang, and T. S. H. Lee, *Phys. Rev.* **C55**, 2024(1997).
3. L. Ya. Glozman and D. O. Riska, *Phys. Reports* **268**, 263(1996);
L. Ya. Glozman, *Nucl. Phys.* **A663**, 103c(2000).
4. N. Isgur and G. Karl, *Phys. Rev.* **D19**, 2653(1979).
5. FI. Stancu, S. Pepin and L. Ya. Glozman, *Phys. Rev* **C56**, 2779(1997).
6. K. Shimizu and L. Ya. Glozman, *Phys. Lett* **B477**, 59(2000).
7. L. R. Dai, Z. Y. Zhang, Y. W. Yu, P. Wang, N-N Interactions in the extended chiral $SU(3)$ quark model, *Nucl. Phys.* **A727**, 321(2003)
8. D. O. Riska and G. E. Grown, *arXiv*, *nucl-th/0005049*, (1999).